

## SARIÇANAK-98

Durum Wheat (*Triticum durum* Desf.)



<b>Registration Year</b>	1998	
<b>Breeding Place and Year</b>	Diyarbakır - 1996	
<b>Applicant Institution</b>	GAP International Agricultural Research and Training Center	
<b>Breeder Institution</b>	GAP International Agricultural Research and Training Center	
<b>Breeding Method</b>	Selection	
<b>Stem and Leaf Characteristics</b>	Medium-height, waxy and strong stemmed, wide-bent leaf	
<b>Spike Structure</b>	Erect, sharp tipped, white and yellow spiked, brown stringy	
<b>Grain Characteristics</b>	Amber colored, vitreous and thousand grain weight 35-45 gr	
<b>Agricultural Characteristics</b>	Spring and medium-early	
<b>Technological Characteristics</b>	<b>1. Test weight (kg/hl)</b>	78-85
	<b>2. Protein (%)</b>	12-16
	<b>3. SDS Value</b>	11-13
	<b>4. Vitreousness (%)</b>	95-100
	<b>5. b yellow colour value</b>	21-23
<b>Yield Situation</b>	Average yield in optimum conditions is 5000 kg/ha, but yield potential can be rise up to 8000 kg/ha.	
<b>Disease Situation</b>	It shows moderately tolerant reaction against yellow rust disease which can be seen from time to time in the region.	
<b>Places Experiments are Performed</b>	Southeastern Anatolia Region's rainfed and irrigated conditions (high yield at irrigation) and Coastal Zones	